

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number
WO 2004/048382 A1

(51) International Patent Classification⁷: C07D 487/04, A61K 31/4162, 31/4709, A61P 35/00

(21) International Application Number:
PCT/US2003/032747

(22) International Filing Date:
10 November 2003 (10.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/428,893 22 November 2002 (22.11.2002) US

(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BEIGHT, Douglas, Wade [US/US]; 3648 South County Road, 600 West, Frankfort, IN 46041 (US). SAWYER, Jason, Scott [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). YINGLING, Jonathan, Michael [US/US]; 10679 Misty Hollow Lane, Fishers, IN 46038 (US).

(74) Agents: SAYLES, Michael, J. et al.; ELI LILLY AND COMPANY, P.O. Box 6288, Indianapolis, IN 46206-6288 (US).

(81) Designated States (national): AE, AG, AL, AM, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, EG, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: QUINOLINYL-PYRROLOPYRAZOLES



Formula II

(57) Abstract: A compound according to formula II and the pharmaceutically acceptable salts thereof and the method of treating cancer in a patient in need thereof by administration of said compound.



— of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

QUINOLINYL-PYRROLOPYRAZOLES

The invention relates to new quionoliny-pyrazole compounds and their use as pharmaceutical agents, in particular their use as TGF-beta signal transduction inhibitors.

5

BACKGROUND OF THE INVENTION

The transforming growth factor-beta (TGF-beta) ("TGF- β ") polypeptides influence growth, differentiation, and gene expression in many cell types. The first polypeptide of this family that was characterized, TGF- β 1, has two identical 112 amino acid subunits that are covalently linked. TGF- β 1 is a highly conserved protein with only a single amino acid difference distinguishing humans from mice. There are two other members of the TGF- β gene family that are expressed in mammals. TGF- β 2 is 71% homologous to TGF- β 1 (de Martin, et al. (1987) EMBO J. 6:3673-3677), whereas 15 TGF- β 3 is 80% homologous to TGF- β 1 (Derynck, et al. (1988) EMBO J 7:3737-3743). The structural characteristics of TGF- β 1 as determined by nuclear magnetic resonance (Archer, et al. (1993) Biochemistry 32:1164-1171) agree with the crystal structure of TGF- β 2 (Daopin, et al. (1992) Science 257:369-374; Schlunegger and Grutter (1992) Nature 358:430-434).

20 There are at least three different extracellular TGF- β receptors, Type I, II and III that are involved in the biological functions of TGF- β 1, - β 2 and - β 3 (For reviews, see Derynck (1994) TIBS 19:548-553 and Massague (1990) Ann. Rev. Cell Biol. 6:597-641). The Type I and Type II receptors are transmembrane serine/threonine kinases, which in the presence of TGF- β form a heteromeric signaling complex (Wrana, et al (1992) 25 Cell 71: 1003-1014).

The mechanism of activation of the heteromeric signaling complex at the cell surface has been elucidated (Wrana, et al. (1994) Nature 370: 341-347). TGF- β first binds the type II receptor that is a constitutively active transmembrane serine/threonine 30 kinase. The type I receptor is subsequently recruited into the complex, phoshorylated at the GS domain and activated to phosphorylate downstream signaling components (e.g. Smad proteins) to initiate the intracellular signaling cascade. A constitutively active type I receptor (T204D mutant) has been shown to effectively transduce TGF- β responses, thus

bypassing the requirement for TGF- β and the type II receptor (Wieser, et al. (1995) EMBO J 14: 2199-2208). Although no signaling function has been discovered for the type III receptor, it does increase TGF- β 2's affinity for the type II receptor making it essentially equipotent with TGF- β 1 and TGF- β 3 (Lopez-Casillas, et al. (1993) Cell
5 73:1435-1444).


Vascular endothelial cells lack the Type III receptor. Instead endothelial cells express a structurally related protein called endoglin (Cheifetz, et al. (1992) J. Biol. Chem. 267:19027-19030), which only binds TGF- β 1 and TGF- β 3 with high affinity. Thus, the relative potency of the TGF- β 's reflects the type of receptors expressed in a cell
10 and organ system. In addition to the regulation of the components in the multi-factorial signaling pathway, the distribution of the synthesis of TGF- β polypeptides also affects physiological function. The distribution of TGF- β 2 and TGF- β 3 is more limited (Derynck, et al. (1988) EMBO J 7:3737-3743) than TGF- β 1, e.g., TGF- β 3 is limited to tissues of mesenchymal origin, whereas TGF- β 1 is present in both tissues of
15 mesenchymal and epithelial origin.

TGF- β 1 is a multifunctional cytokine critical for tissue repair. High concentrations of TGF- β 1 are delivered to the site of injury by platelet granules (Assoian and Sporn (1986) J. Cell Biol. 102:1217-1223). TGF- β 1 initiates a series of events that promote healing including chemo taxis of cells such as leukocytes, monocytes and
20 fibroblasts, and regulation of growth factors and cytokines involved in angiogenesis, cell division associated with tissue repair and inflammatory responses. TGF- β 1 also stimulates the synthesis of extracellular matrix components (Roberts, et al. (1986) Proc. Natl. Acad. Sci. USA 83:4167-4171; Sporn, et al. (1983) Science 219:1329-1330; Massague (1987) Cell 49:437-438) and most importantly for understanding the
25 pathophysiology of TGF- β 1, TGF- β 1 autoregulates its own synthesis (Kim, et al. (1989) J. Biol. Chem. 264:7041-7045).

The compounds disclosed herein may also exhibit other kinase activity, such as p38 kinase inhibition and/or KDR (VEGFR2) kinase inhibition. Assays to determine such kinase activity are known in the art and one skilled in the art would be able to test the
30 disclosed compounds for such activity.

SUMMARY OF THE INVENTION

The disclosed invention also relates to the select compound of Formula II:



Formula II

2-(6-methyl-pyridin-2-yl)-3-[6-amido-quinolin-4-yl]-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole

and the pharmaceutically acceptable salts thereof.

The compound above is generically disclosed and claimed in PCT patent application PCT/US02/11884, filed 13 May 2002, which claims priority from U.S. patent application U.S.S.N. 60/293,464, filed 24 May 2001, and incorporated herein by reference. The above compound has been selected for having a surprisingly superior toxicology profile over the compounds specifically disclosed in application cited above.

DETAILED DESCRIPTION OF THE INVENTION

The term "effective amount" as used in "an effective amount of a compound of Formula I," for example, refers to an amount of a compound of the present invention that is capable of inhibiting TGF-beta.

The term μM refers to micromolar.

The general chemical terms used herein have their usual meanings.

The following abbreviations are used throughout the synthesis Schemes and Examples:

DMF refers to dimethyl formamide

THF refers to tetrahydrofuran


Ms refers to mesyl which is methylsulfonyl

THP refers to tetrahydropyran

-4-


The compounds disclosed herein can be made according to the following schemes and examples. The examples should in no way be understood to be limiting in any way as to how the compounds may be made.

5 The following scheme illustrates the preparation of the compound of Formula I.



-5-

The following scheme illustrates the preparation of the compound of Formula II.



5

The following examples further illustrate the preparation of the compounds of this invention as shown schematically in Schemes I and II.

Example 1**Preparation of 7-(2-morpholin-4-yl-ethoxy)-4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinoline****A. Preparation of 4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-7-[2-(tetrahydropyran-2-yloxy)ethoxy]quinoline**

Heat 4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinolin-7-ol (376 mg, 1.146 mmol), cesium carbonate (826 mg, 2.54 mmol), and 2-(2-bromoethoxy)tetrahydro-2H-pyran (380 μ L, 2.52 mmol) in DMF (5 mL) at 120 °C for 4 hours. Quench the reaction with saturated sodium chloride and then extract with chloroform. Dry the organic layer over sodium sulfate and concentrate in vacuo. Purify the reaction mixture on a silica gel column eluting with dichloromethane to 10% methanol in dichloromethane to give the desired subtitled intermediate as a yellow oil (424 mg, 81%). MS ES⁺ m/e 457.0 (M+1).

B. Preparation of 2-[4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinolin-7-yloxy]-ethanol


Heat a solution of 4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-7-[2-(tetrahydropyran-2-yloxy)ethoxy]quinoline (421 mg, 0.92 mmol) in acetic acid:tetrahydrofuran: water (4:2:1) (20 mL). Remove the solvent in vacuo and recover the residue with chloroform:isopropyl (3:1). Wash the organic layer with saturated sodium bicarbonate and dry over sodium sulfate. Concentrate in vacuo. The residue will be pure enough for the next step in the scheme (425 mg, 100%). MS ES⁺ m/e 373.1 (M+1).

C. Preparation of methanesulfonic acid 2-[4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinolin-7-yloxy]-ethyl ester

Stir a solution of 2-[4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinolin-7-yloxy]-ethanol (293 mg, 0.78 mmol) and methane sulfonyl chloride (68 μ L, 0.81 ml) in dried pyridine (5 mL) for 2 hours. Remove the pyridine in vacuo and recover the residue with chloroform. Wash the organic layer with saturated sodium bicarbonate and dry over sodium sulfate to give the desired subtitled intermediate as a white foam (425 mg, 100%). MS ES⁺ m/e 451.1 (M+1).

-7-

D. Preparation of 7-(2-morpholin-4-yl-ethoxy)-4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinoline



5 Heat methanesulfonic acid 2-[4-(2-pyridin-2-yl-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)-quinolin-7-yloxy]-ethyl ester (87 mg, 0.19 mmol) with morpholine (1 mL) at 50 °C for 4 hours. Remove the morpholine in vacuo and then extract the product with isopropyl alcohol:chloroform (1:3). Wash the organic layer with sodium chloride
10 and dry over sodium sulfate. Concentrate in vacuo to give the desired title product as a slight yellow solid (83 mg, 100%). MS ES⁺ m/e 442.0 (M+1).

EXAMPLE 2

15

Preparation of

2-(6-methyl-pyridin-2-yl)-3-[6-amido-quinolin-4-yl]-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole

20 **A. Preparation of 6-bromo-4-methyl-quinoline**

Stir a solution of 4-bromo-phenylamine (1 eq), in 1,4-dioxane and cool to approximately 12 °C. Slowly add sulfuric acid (2 eq) and heat at reflux. Add methylvinyl ketone (1.5 eq) dropwise into the refluxing solution. Heat the solution for 1
25 hour after addition is complete. Evaporate the reaction solution to dryness and dissolve in methylene chloride. Adjust the solution to pH 8 with 1 M sodium carbonate and extract three times with water. Chromatograph the residue on SiO₂ (70/30 hexane/ethyl acetate) to obtain the desired subtitled intermediate.

MS ES⁺ m/e = 158.2 (M+1).

30

B. Preparation of 6-methyl-pyridine-2-carboxylic acid methyl ester

Suspend 6-methyl-pyridine-2-carboxylic acid (10 g, 72.9 mmol) in methylene chloride (200 mL). Cool to 0 °C. Add methanol (10 mL), 4-dimethylaminopyridine (11.6 g, 94.8 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (18.2 g, 94.8 mmol). Stir the mixture at room temperature for 6 hours, wash with water and brine, and dry over sodium sulfate. Filter the mixture and concentrate in vacuo. Chromatograph the residue on SiO₂ (50% ethyl acetate/hexanes) to obtain the desired subtitled intermediate, 9.66 g (92%), as a colorless liquid.

¹H NMR (CDCl₃) δ 7.93-7.88 (m, 1H), 7.75-7.7 (m, 1H), 7.35-7.3 (m, 1H), 4.00 (s, 3H), 2.60 (s, 3H).

C. Preparation of 2-(6-bromo-quinolin-4-yl)-1-(6-methyl-pyridin-2-yl)-ethanone

Dissolve 6-bromo-4-methyl-quinoline (38.5 g, 153 mmol) in 600 mL dry THF. Cool to -70° C and treat with the dropwise addition of 0.5 M potassium hexamethyldisilazane (KN(SiMe₃)₂) (400 mL, 200 mmol) over 2 hours while keeping the temperature below -65 °C. Stir the resultant solution at -70°C for 1 hour and add a solution of 6-methylpyridine-2-carboxylic acid methyl ester (27.2, 180 mmol) in 100 mL dry THF dropwise over 15 minutes. During the addition, the mixture will turn from dark red to pea-green and form a precipitate. Stir the mixture at -70°C over 2 hours then allow it to warm to ambient temperature with stirring for 5 hours. Cool the mixture then quench with 12 N HCl to pH=1. Raise the pH to 9 with solid potassium carbonate. Decant the solution from the solids and extract twice with 200 mL ethyl acetate. Combine the organic extracts, wash with water and dry over potassium carbonate. Stir the solids in 200 mL water and 200 mL ethyl acetate and treat with additional potassium carbonate. Separate the organic portion and dry with the previous ethyl acetate extracts. Concentrate the solution in vacuo to a dark oil. Pass the oil through a 300 mL silica plug with methylene chloride then ethyl acetate. Combine the appropriate fractions and concentrate in vacuo to yield an amber oil. Rinse the oil down the sides of the flask with methylene chloride then dilute with hexane while swirling the flask to yield 38.5 g (73.8 %) of the desired subtitled intermediate as a yellow solid.

MS ES+ = 341 (M+1).

D. Preparation of 1-[2-(6-bromo-quinolin-4-yl)-1-(6-methyl-pyridin-2-yl)-ethylideneamino]-pyrrolidin-2-one

Stir a mixture of 2-(6-bromo-quinolin-4-yl)-1-(6-methyl-pyridin-2-yl)-ethanone
5 (38.5 g, 113 mmol) and 1-aminopyrrolidinone hydrochloride (20 g, 147 mmol) in 115 mL pyridine at ambient temperature for 10 hours. Add about 50 g 4 Å unactivated sieves. Continue stirring an additional 13 h and add 10-15 g silica and filter the mixture through a 50 g silica plug. Elute the silica plug with 3 L ethyl acetate. Combine the filtrates and concentrate in vacuo. Collect the hydrazone precipitate by
10 filtration and suction dry to yield 33.3 g (69.7%) of the desired subtitled intermediate as an off-white solid.
MS ES⁺ = 423 (M+1).

E. Preparation of 6-bromo-4-[2-(6-methyl-pyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl]-quinoline

To a mixture of (1.2 eq.) cesium carbonate and 1-[2-(6-bromo-quinolin-4-yl)-1-(6-methyl-pyridin-2-yl)-ethylideneamino]-pyrrolidin-2-one (33.3 g, 78.7 mmol) add
300 mL dry N,N-dimethylformamide. Stir the mixture 20 hours at 100°C. The
20 mixture may turn dark during the reaction. Remove the N,N-dimethylformamide in vacuo. Partition the residue between water and methylene chloride. Extract the aqueous portion with additional methylene chloride. Filter the organic solutions through a 300 mL silica plug, eluting with 1.5 L methylene chloride, 1.5 L ethyl acetate and 1.5 L acetone. Combine the appropriate fractions and concentrate in
25 vacuo. Collect the resulting precipitate by filtration to yield 22.7 g (71.2%) of the desired subtitled intermediate as an off-white solid.
MS ES⁺ = 405 (M+1).

F. Preparation of 4-[2-(6-methyl-pyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl]-quinoline-6-carboxylic acid methyl ester


Add 6-bromo-4-[2-(6-methyl-pyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl]-quinoline (22.7 g, 45 mmol) to a mixture of sodium acetate (19 g, 230 mmol) and the palladium catalyst [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with
35

-10-

dichloromethane (1:1) (850 mg, 1.04 mmol) in 130 mL methanol. Place the mixture under 50 psi carbon monoxide atmosphere and stir while warming to 90° C over 1 hour and with constant charging with additional carbon monoxide. Allow the mixture to cool over 8 hours, recharge again with carbon monoxide and heat to 90 °C. The pressure may rise to about 75 PSI. The reaction is complete in about an hour when the pressure is stable and tlc (1:1 toluene/acetone) shows no remaining bromide. Partition the mixture between methylene chloride (600 mL) and water (1 L). Extract the aqueous portion with an additional portion of methylene chloride (400 mL.) Filter the organic solution through a 300 mL silica plug and wash with 500 mL methylene chloride, 1200 mL ethyl acetate and 1500 mL acetone. Discard the acetone portion. Combine appropriate fractions and concentrate to yield 18.8 g (87.4%) of the desired subtitled intermediate as a pink powder.

MS ES⁺ = 385 (M+1).

15 **G. Preparation of 2-(6-methyl-pyridin-2-yl)-3-[6-amido-quinolin-4-yl]-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole**



20 Warm a mixture of 4-[2-(6-methyl-pyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl]-quinoline-6-carboxylic acid methyl ester in 60 mL 7 N ammonia in methanol to 90 °C in a stainless steel pressure vessel for 66 hours. The pressure will rise to about 80 PSI. Maintain the pressure for the duration of the reaction. Cool the vessel and concentrate the brown mixture in vacuo. Purify the residual solid on two 12 g Redi-

25 Pak cartridges coupled in series eluting with acetone. Combine appropriate fractions and concentrate in vacuo. Suspend the resulting nearly white solid in methylene chloride, dilute with hexane, and filter. The collected off-white solid yields 1.104 g (63.8%) of the desired title product.

MS ES⁺ = 370 (M+1).

-11-

The compounds disclosed herein were tested by the following protocols for TGF- β inhibition, as described below in the protocol description.

TGF- β RECEPTOR I PURIFICATION AND *IN VITRO* KINASE

5

REACTIONS

For TGF- β Type I (RIT204D) Receptors:

The 6X-HIS tagged cytoplasmic kinase domain of each receptor was expressed and purified from Sf9 insect cell lysates as briefly described below:

10 Cell pellets after 48-72 hours of infection were lysed in lysis buffer (LB: 50 mM Tris pH 7.5, 150 mM NaCl, 50 mM NaF, 0.5% NP40 with freshly added 20 mM β -mercaptoethanol, 10 mM imidazole, 1 mM PMSF, 1X EDTA-free Complete Protease Inhibitor(Boehringer Mannheim).

Cell lysates were clarified by centrifugation and 0.45 μ M filtered prior to
15 purification by Ni/NTA affinity chromatography (Qiagen).

Chromatography Protocol:

Equilibrate with 10 CV of LB, load sample, wash with 10 CV RIPA buffer (50
20 mM Tris pH 7.5, 150 mM NaCl, 1% NP40, 1mM EDTA, 0.25% sodium deoxycholate, added fresh 20 mM β -mercaptoethanol, 1 mM PMSF), wash with 10 CV LB, wash with 10 CV 1X KB (50 mM Tris pH 7.5, 150 mM NaCl, 4 mM $MgCl_2$, 1 mM NaF, 2 mM β -mercaptoethanol), elute with a linear gradient of 1X KB containing 200 mM Imidazole.

Both enzymes were approximately 90% pure and had autophosphorylation
25 activity.

Reactions: 170-200 nM enzyme in 1X KB, compound dilution series in 1X KB/16% DMSO (20 μ M to 1 nM final concentration with 4% DMSO final concentration), reactions started by adding ATP mix (4 μ M ATP/1 μ Ci ^{33}P - γ -ATP final concentrations) in 1X KB.

30 Reactions were incubated at 30 $^{\circ}C$ for 1 hour. Reactions were stopped and quantitated using standard TCA/BSA precipitation onto Millipore FB glass fiber filter plates and by liquid scintillation counting on a MicroBeta JET.

The compounds disclosed herein inhibit the TGF- β Type I (RIT204D) receptor kinase domain with IC₅₀ values <20 μ M, while exhibiting less toxicity *in vivo* than structurally related compounds as disclosed in PCT patent application PCT/US02/11884 identified above.

5 Conditions "characterized by enhanced TGF- β activity" include those wherein TGF- β synthesis is stimulated so that TGF- β is present at increased levels or wherein TGF- β latent protein is undesirably activated or converted to active TGF- β protein or wherein TGF- β receptors are upregulated or wherein the TGF- β protein shows enhanced binding to cells or extracellular matrix in the location of the disease. Thus, in either case
10 "enhanced activity" refers to any condition wherein the biological activity of TGF- β is undesirably high, regardless of the cause.

A number of diseases have been associated with TGF- β 1 over production. Inhibitors of TGF- β intracellular signaling pathway are useful treatments for fibroproliferative diseases. Specifically, fibroproliferative diseases include kidney
15 disorders associated with unregulated TGF- β activity and excessive fibrosis including glomerulonephritis (GN), such as mesangial proliferative GN, immune GN, and crescentic GN. Other renal conditions include diabetic nephropathy, renal interstitial fibrosis, renal fibrosis in transplant patients receiving cyclosporin, and HIV-associated nephropathy. Collagen vascular disorders include progressive systemic sclerosis,
20 polymyositis, scleroderma, dermatomyositis, eosinophilic fascitis, morphea, or those associated with the occurrence of Raynaud's syndrome. Lung fibroses resulting from excessive TGF- β activity include adult respiratory distress syndrome, idiopathic pulmonary fibrosis, and interstitial pulmonary fibrosis often associated with autoimmune disorders, such as systemic lupus erythematosus and scleroderma, chemical contact, or
25 allergies. Another autoimmune disorder associated with fibroproliferative characteristics is rheumatoid arthritis.

Eye diseases associated with a fibroproliferative condition include retinal reattachment surgery accompanying proliferative vitreoretinopathy, cataract extraction with intraocular lens implantation, and post glaucoma drainage surgery are associated
30 with TGF- β 1 overproduction.

Fibrotic diseases associated with TGF- β 1 overproduction can be divided into chronic conditions such as fibrosis of the kidney, lung and liver and more acute conditions such as dermal scarring and restenosis (Chamberlain, J. Cardiovascular Drug Reviews, 19(4):329-344). Synthesis and secretion of TGF- β 1 by tumor cells can also
5 lead to immune suppression such as seen in patients with aggressive brain or breast tumors (Arteaga, et al. (1993) J. Clin. Invest. 92:2569-2576). The course of Leishmanial infection in mice is drastically altered by TGF- β 1 (Barral-Netto, et al. (1992) Science 257:545-547). TGF- β 1 exacerbated the disease, whereas TGF- β 1 antibodies halted the progression of the disease in genetically susceptible mice. Genetically resistant mice
10 became susceptible to Leishmanial infection upon administration of TGF- β 1.

The profound effects of TGF- β 1 on extracellular matrix deposition have been reviewed (Rocco and Ziyadeh (1991) in Contemporary Issues in Nephrology v.23, Hormones, autocoids and the kidney. ed. Jay Stein, Churchill Livingston, New York pp.391-410; Roberts, et al. (1988) Rec. Prog. Hormone Res. 44:157-197) and include the
15 stimulation of the synthesis and the inhibition of degradation of extracellular matrix components. Since the structure and filtration properties of the glomerulus are largely determined by the extracellular matrix composition of the mesangium and glomerular membrane, it is not surprising that TGF- β 1 has profound effects on the kidney. The accumulation of mesangial matrix in proliferative glomerulonephritis (Border, et al.
20 (1990) Kidney Int. 37:689-695) and diabetic nephropathy (Mauer, et al. (1984) J. Clin. Invest. 74:1143-1155) are clear and dominant pathological features of the diseases. TGF- β 1 levels are elevated in human diabetic glomerulosclerosis (advanced neuropathy) (Yamamoto, et al. (1993) Proc. Natl. Acad. Sci. 90:1814-1818). TGF- β 1 is an important mediator in the genesis of renal fibrosis in a number of animal models (Phan, et al. (1990)
25 Kidney Int. 37:426; Okuda, et al. (1990) J. Clin. Invest. 86:453). Suppression of experimentally induced glomerulonephritis in rats has been demonstrated by antiserum against TGF- β 1 (Border, et al. (1990) Nature 346:371) and by an extracellular matrix protein, decorin, which can bind TGF- β 1 (Border, et al. (1992) Nature 360:361-363).

Too much TGF- β 1 leads to dermal scar-tissue formation. Neutralizing TGF- β 1
30 antibodies injected into the margins of healing wounds in rats have been shown to inhibit scarring without interfering with the rate of wound healing or the tensile strength of the

wound (Shah, et al. (1992) Lancet 339:213-214). At the same time there was reduced angiogenesis, reduced number of macrophages and monocytes in the wound, and a reduced amount of disorganized collagen fiber deposition in the scar tissue.

TGF- β 1 may be a factor in the progressive thickening of the arterial wall which results from the proliferation of smooth muscle cells and deposition of extracellular matrix in the artery after balloon angioplasty. The diameter of the restenosed artery may be reduced 90% by this thickening, and since most of the reduction in diameter is due to extracellular matrix rather than smooth muscle cell bodies, it may be possible to open these vessels to 50% simply by reducing extensive extracellular matrix deposition. In uninjured pig arteries transfected in vivo with a TGF- β 1 gene, TGF- β 1 gene expression was associated with both extracellular matrix synthesis and hyperplasia (Nabel, et al. (1993) Proc. Natl. Acad. Sci. USA 90:10759-10763). The TGF- β 1 induced hyperplasia was not as extensive as that induced with PDGF-BB, but the extracellular matrix was more extensive with TGF- β 1 transfectants. No extracellular matrix deposition was associated with FGF-1 (a secreted form of FGF) induced hyperplasia in this gene transfer pig model (Nabel (1993) Nature 362:844-846).

There are several types of cancer where TGF- β 1 produced by the tumor may be deleterious. MATLyLu rat prostate cancer cells (Steiner and Barrack (1992) Mol. Endocrinol 6:15-25) and MCF-7 human breast cancer cells (Arteaga, et al. (1993) Cell Growth and Differ. 4:193-201) became more tumorigenic and metastatic after transfection with a vector expressing the mouse TGF- β 1. TGF- β 1 has been associated with angiogenesis, metastasis and poor prognosis in human prostate and advanced gastric cancer (Wikstrom, P., et al. (1998) Prostate 37: 19-29; Saito, H. et al. (1999) Cancer 86: 1455-1462). In breast cancer, poor prognosis is associated with elevated TGF- β (Dickson, et al. (1987) Proc. Natl. Acad. Sci. USA 84:837-841; Kasid, et al. (1987) Cancer Res. 47:5733-5738; Daly, et al. (1990) J. Cell Biochem. 43:199-211; Barrett-Lee, et al. (1990) Br. J Cancer 61:612-617; King, et al. (1989) J. Steroid Biochem. 34:133-138; Welch, et al. (1990) Proc. Natl. Acad. Sci. USA 87:7678-7682; Walker, et al. (1992) Eur. J. Cancer 238:641-644) and induction of TGF- β 1 by tamoxifen treatment (Butta, et al. (1992) Cancer Res. 52:4261- 4264) has been associated with failure of tamoxifen treatment for breast cancer (Thompson, et al. (1991) Br. J. Cancer 63:609-614). Anti

TGF- β 1 antibodies inhibit the growth of MDA-231 human breast cancer cells in athymic mice (Arteaga, et al. (1993) J. Clin. Invest. 92:2569-2576), a treatment which is correlated with an increase in spleen natural killer cell activity. CHO cells transfected with latent TGF- β 1 also showed decreased NK activity and increased tumor growth in nude mice (Wallick, et al. (1990) J. Exp. Med. 172:1777-1784). Thus, TGF- β secreted by breast tumors may cause an endocrine immune suppression. High plasma concentrations of TGF- β 1 have been shown to indicate poor prognosis for advanced breast cancer patients (Anscher, et al. (1993) N. Engl. J. Med. 328:1592-1598). Patients with high circulating TGF- β before high dose chemotherapy and autologous bone marrow transplantation are at high risk for hepatic veno-occlusive disease (15-50% of all patients with a mortality rate up to 50%) and idiopathic interstitial pneumonitis (40-60% of all patients). The implication of these findings is 1) that elevated plasma levels of TGF- β 1 can be used to identify at risk patients and 2) that reduction of TGF- β 1 could decrease the morbidity and mortality of these common treatments for breast cancer patients.

Many malignant cells secrete transforming growth factor- β (TGF- β), a potent immunosuppressant, suggesting that TGF- β production may represent a significant tumor escape mechanism from host immunosurveillance. Establishment of a leukocyte subpopulation with disrupted TGF- β signaling in the tumor-bearing host offers a potential means for immunotherapy of cancer. A transgenic animal model with disrupted TGF- β signaling in T cells is capable of eradicating a normally lethal TGF- β over expressing lymphoma tumor, EL4 (Gorelik and Flavell, (2001) Nature Medicine 7(10): 1118-1122). Down regulation of TGF- β secretion in tumor cells results in restoration of immunogenicity in the host, while T-cell insensitivity to TGF- β results in accelerated differentiation and autoimmunity, elements of which may be required in order to combat self-antigen-expressing tumors in a tolerized host. The immunosuppressive effects of TGF- β have also been implicated in a subpopulation of HIV patients with lower than predicted immune response based on their CD4/CD8 T cell counts (Garba, et al. J. Immunology (2002) 168: 2247-2254). A TGF- β neutralizing antibody was capable of reversing the effect in culture, indicating that TGF- β signaling inhibitors may have utility in reversing the immune suppression present in this subset of HIV patients.

During the earliest stages of carcinogenesis, TGF- β 1 can act as a potent tumor suppressor and may mediate the actions of some chemopreventive agents. However, at some point during the development and progression of malignant neoplasms, tumor cells appear to escape from TGF- β -dependent growth inhibition in parallel with the appearance of bioactive TGF- β in the microenvironment. The dual tumor suppression/tumor promotion roles of TGF- β have been most clearly elucidated in a transgenic system over expressing TGF- β in keratinocytes. While the transgenics were more resistant to formation of benign skin lesions, the rate of metastatic conversion in the transgenics was dramatically increased (Cui, et al (1996) Cell 86(4):531-42). The production of TGF- β 1 by malignant cells in primary tumors appears to increase with advancing stages of tumor progression. Studies in many of the major epithelial cancers suggest that the increased production of TGF- β by human cancers occurs as a relatively late event during tumor progression. Further, this tumor-associated TGF- β provides the tumor cells with a selective advantage and promotes tumor progression. The effects of TGF- β on cell/cell and cell/stroma interactions result in a greater propensity for invasion and metastasis. Tumor-associated TGF- β may allow tumor cells to escape from immune surveillance since it is a potent inhibitor of the clonal expansion of activated lymphocytes. TGF- β has also been shown to inhibit the production of angiostatin. Cancer therapeutic modalities such as radiation therapy and chemotherapy induce the production of activated TGF- β in the tumor, thereby selecting outgrowth of malignant cells that are resistant to TGF- β growth inhibitory effects. Thus, these anticancer treatments increase the risk and hasten the development of tumors with enhanced growth and invasiveness. In this situation, agents targeting TGF- β -mediated signal transduction might be a very effective therapeutic strategy. The resistance of tumor cells to TGF- β has been shown to negate much of the cytotoxic effects of radiation therapy and chemotherapy and the treatment-dependent activation of TGF- β in the stroma may even be detrimental as it can make the microenvironment more conducive to tumor progression and contributes to tissue damage leading to fibrosis. The development of a TGF- β signal transduction inhibitors is likely to benefit the treatment of progressed cancer alone and in combination with other therapies.

-17-

The compounds are useful for the treatment of cancer and other disease states influenced by TGF- β by inhibiting TGF- β in a patient in need thereof by administering said compound(s) to said patient. TGF- β would also be useful against atherosclerosis (T.A. McCaffrey: TGF- β s and TGF- β Receptors in Atherosclerosis: Cytokine and Growth Factor Reviews 2000, 11, 103-114) and Alzheimer's (Masliah, E.; Ho, G.; Wyss-Coray, T.: Functional Role of TGF- β in Alzheimer's Disease Microvascular Injury: Lessons from Transgenic Mice: Neurochemistry International 2001, 39, 393-400) diseases.

PHARMACEUTICAL COMPOSITIONS

The compositions of the present invention are therapeutically effective amounts of the TGF- β antagonists, noted above. The composition may be formulated with common excipients, diluents or carriers, and compressed into tablets, or formulated elixirs or solutions for convenient oral administration or administered by intramuscular intravenous routes. The compounds can be administered transdermally and maybe formulated as sustained release dosage forms and the like.

The method of treating a human patient according to the present invention includes administration of the TGF- β antagonists. The TGF- β antagonists are formulated into formulations which may be administered by the oral and rectal routes, topically, parenterally, e.g., by injection and by continuous or discontinuous intra-arterial infusion, in the form of, for example, tablets, lozenges, sublingual tablets, sachets, cachets, elixirs, gels, suspensions, aerosols, ointments, for example, containing from 1 to 10% by weight of the active compound in a suitable base, soft and hard gelatin capsules, suppositories, injectable solutions and suspensions in physiologically acceptable media, and sterile packaged powders adsorbed onto a support material for making injectable solutions. Advantageously for this purpose, compositions may be provided in dosage unit form, preferably each dosage unit containing from about 5 to about 500 mg (from about 5 to 50 mg in the case of parenteral or inhalation administration, and from about 25 to 500 mg in the case of oral or rectal administration) the compounds. Dosages from about 0.5 to about 300 mg/kg per day, preferably 0.5 to 20 mg/kg, of active ingredient may be administered although it will, of course, readily be understood that the amount of the compound actually to be administered will be determined by a physician, in the light of all the relevant circumstances including the condition to be treated, the choice of compound to


be administered and the choice of route of administration and therefore the above preferred dosage range is not intended to limit the scope of the present invention in any way.

The formulations useful for separate administration of the TGF- β antagonists will normally consist of at least one compound selected from the compounds specified herein mixed with a carrier, or diluted by a carrier, or enclosed or encapsulated by an ingestible carrier in the form of a capsule, sachet, cachet, paper or other container or by a disposable container such as an ampoule. A carrier or diluent may be a solid, semi-solid or liquid material which serves as a vehicle, excipient or medium for the active therapeutic substance. Some examples of the diluents or carrier which may be employed in the pharmaceutical compositions of the present invention are lactose, dextrose, sucrose, sorbitol, mannitol, propylene glycol, liquid paraffin, white soft paraffin, kaolin, fumed silicon dioxide, microcrystalline cellulose, calcium silicate, silica, polyvinylpyrrolidone, cetostearyl alcohol, starch, modified starches, gum acacia, calcium phosphate, cocoa butter, ethoxylated esters, oil of theobroma, arachis oil, alginates, tragacanth, gelatin, syrup, methyl cellulose, polyoxyethylene sorbitan monolaurate, ethyl lactate, methyl and propyl hydroxybenzoate, sorbitan trioleate, sorbitan sesquioleate and oleyl alcohol and propellants such as trichloromonofluoromethane, dichlorodifluoromethane and dichlorotetrafluoroethane. In the case of tablets, a lubricant may be incorporated to prevent sticking and binding of the powdered ingredients in the dies and on the punch of the tableting machine. For such purpose there may be employed for instance aluminum, magnesium or calcium stearates, talc or mineral oil.

Preferred pharmaceutical forms of the present invention are capsules, tablets, suppositories, injectable solutions, creams and ointments. Especially preferred are formulations for inhalation application, such as an aerosol, for injection, and for oral ingestion.

WE CLAIM:

1. A compound according to formula II



Formula II

and the pharmaceutically acceptable salts thereof.

2. The compound which is 2-(6-methyl-pyridin-2-yl)-3-[6-amido-quinolin-4-yl]-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole and the pharmaceutically acceptable salts thereof.

3. A pharmaceutical formulation comprising a compound according to Claim 1 or the pharmaceutically acceptable salt, ester or prodrug thereof together with a pharmaceutically acceptable diluent or carrier.

4. The method of treating cancer which comprises administering to a patient in need thereof a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt, ester or prodrug thereof.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 03/32747

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D487/04 A61K31/4162 A61K31/4709 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 02 094833 A (CIAPETTI PAOLA ;BEIGHT DOUGLAS WADE (US); GODFREY ALEXANDER GLENN) 28 November 2002 (2002-11-28) page 193; examples 282-285	1-4
A	EP 0 531 901 A (FUJISAWA PHARMACEUTICAL CO) 17 March 1993 (1993-03-17) claims; examples	1-4

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 March 2004

Date of mailing of the international search report

31/03/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fazzi, R

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 03/32747

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 4 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International patent family members

International Application No
PCT/US 03/32747

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 02094833	A	28-11-2002	CA 2446820 A1	28-11-2002
			EP 1397364 A1	17-03-2004
			NO 20035193 A	21-11-2003
			WO 02094833 A1	28-11-2002
<hr/>				
EP 0531901	A	17-03-1993	AU 2280592 A	11-03-1993
			CA 2077732 A1	10-03-1993
			CN 1070404 A	31-03-1993
			EP 0531901 A2	17-03-1993
			HU 65204 A2	02-05-1994
			JP 7252256 A	03-10-1995
			JP 2057962 C	10-06-1996
			JP 6287188 A	11-10-1994
			JP 7088386 B	27-09-1995
			MX 9205145 A1	01-04-1993
			US 5478827 A	26-12-1995
			US 5624931 A	29-04-1997
			US 5356897 A	18-10-1994
			ZA 9206417 A	15-03-1993
			CN 1075965 A	08-09-1993
<hr/>				